

The BFKL-Regge Expansion for the Proton Structure Function at Small x

Vladimir R. Zoller

*Institute for Theoretical and Experimental Physics
117218 Moscow, Russia*¹

Abstract. We report an evaluation of subleading eigenvalues and eigenfunctions of the BFKL equation in the color dipole representation with a running gauge coupling. We present an expansion of the small- x proton structure function $F_{2p}(x, Q^2)$ in terms of the rightmost BFKL-singularities. The BFKL-Regge phenomenology of DIS structure functions is developed which is shown to provide remarkably good description of the data on $F_{2p}(x, Q^2)$ from E665 to HERA.

Introduction

In this talk I address the issue of extrapolation of the proton structure functions to a domain of very small x from the attainable kinematical region of x and Q^2 . So, this is an old problem of predicting the future from the known past. The standard predictor is the GLDAP evolution equation [1]. However, the art of predicting is difficult. The numerous pre-HERA GLDAP fits to the proton structure functions are equally good for large- x -data but seldom come close to the new data points at smaller x . The lack of predictive power is not surprising and we comment on this point below. It is well known that the GLDAP evolution breaks down at small x and is superseded by the $\log(1/x)$ -BFKL-evolution [2]. Contrary to the GLDAP approach, the BFKL evolution predicts uniquely the proton structure functions at an arbitrarily small x from the input at a starting point $x = x_0$. DIS structure functions at asymptotically large $1/x$ are dominated by the rightmost Regge-singularity with the intercept Δ_0 [3] and

$$F_{2p}(x, Q^2) = F_2^{(0)}(Q^2) \left(\frac{1}{x}\right)^{\Delta_0}. \quad (1)$$

¹⁾ Talk at 5th International Workshop on Deep Inelastic Scattering and QCD (DIS'97), Chicago, April 1997

At moderate x , however, the subleading contributions to $F_{2p}(x, Q^2)$ with smaller intercepts $\Delta_1, \Delta_2, \dots$ can not be neglected. The task of my talk is to present an evaluation of the intercepts Δ_n , the pomeron trajectory slopes α'_n and the corresponding structure functions $F_2^n(Q^2)$, to arrive finally at the representation

$$F_{2p}(x, Q^2) = \sum_n A_n F_2^{(n)}(Q^2) \left(\frac{x_0}{x} \right)^{\Delta_n}. \quad (2)$$

I conclude with the expansion of the proton structure function $F_2(x, Q^2)$ in terms of the three rightmost BFKL singularities. Such a three pole approximation seems to exhaust the existing experimental data thus providing a reliable basis for the BFKL-Regge phenomenology of diffractive DIS.

The BFKL Eigenvalue Problem in the Dipole Picture

The virtual photo-absorption cross section $\sigma(r, x)$, where r is the color dipole size, satisfies the BFKL equation [4] (hereafter $\xi = \log(1/x)$):

$$\frac{\partial \sigma(\xi, r)}{\partial \xi} = \mathcal{K} \otimes \sigma(r, x) \quad (3)$$

with the kernel $\mathcal{K}$ which involves the running gauge coupling and the infrared cutoff- the correlation radius R_c of perturbative gluons. We look for the solution with the Regge behavior

$$\sigma_n(r, x) = \sigma_n(r) \left(\frac{1}{x} \right)^{\Delta_n}. \quad (4)$$

Then the eigenfunctions $\sigma_n(r)$ and the eigenvalues Δ_n are determined from

$$\mathcal{K} \otimes \sigma_n(r) = \Delta_n \sigma_n(r). \quad (5)$$

The short-distance asymptotics of the eigenfunctions is known in the analytic form [5]

$$\sigma_n(r) = r^2 \left[\frac{1}{\alpha_S(r)} \right]^{\gamma_n - 1}, \quad (6)$$

where $\gamma_n \Delta_n = 4/3$. At large distances, $r \gg R_c$,

$$\sigma_n(r) \equiv \bar{\sigma}_n = \text{const} \quad (7)$$

due to the finite R_c . Another useful clue is that the leading eigenfunction $\sigma_0(r)$ is node free and the n -th subleading solution must have n nodes. Then a practical approach to the eigenvalue problem is a variational procedure [6] applied to a class of n -node polynomials $\mathcal{P}_n(z)$ in a variable $z \sim [1/\alpha_S(r)]^\gamma$.

Results and Discussion

The BFKL equation with running coupling and infrared cutoff has a discrete spectrum. The eigenvalues Δ_n obtained by the variational method for $n = 0, 1, 2, 3, \dots$ are as follows

$$\Delta_n = 0.40, 0.220, 0.148, 0.111, 0.088, 0.073, 0.063, \dots \quad (8)$$

To an accuracy better than 10% the above series follows the law

$$\Delta_n = \frac{\Delta_0}{(n+1)} \quad (9)$$

derived by Lipatov [3] from quasi-classical considerations.

The BFKL eigenfunctions are represented (Figure 1) in term of the quantity $\sigma_n(r)/r$ which to a crude approximation is similar to Lipatov's quasi-classical eigenfunctions, which are $\mathcal{E}_n(r) \sim \cos[\phi(r)]$ for $n \gg 1$ [3].

Once $\sigma_n(r)$ is known, the pomeron trajectory slope α'_n can readily be derived. In the notations of ref. [7]

$$\alpha'_n = \frac{3}{32\pi R_c^2} \int d\rho^2 \rho^2 \alpha_S(\rho) K_1^2(\rho/R_c) [1 + \sigma_n(\rho)/\bar{\sigma}_n] , \quad (10)$$

where (see eq.7)

$$\bar{\sigma}_n = \frac{3}{2\pi R_c^2} \int d\rho^2 \alpha_S(\rho) K_1^2(\rho/R_c) \sigma_n(\rho) . \quad (11)$$




FIGURE 1. BFKL eigenfunctions.

Numerically, $\alpha'_0 = 0.072 \text{ GeV}^{-2}$, $\alpha'_1 = 0.066 \text{ GeV}^{-2}$, $\alpha'_2 = 0.062 \text{ GeV}^{-2}$, $\alpha'_3 = 0.060 \text{ GeV}^{-2}$.

The color dipole factorization relates the dipole cross sections with the structure functions $F_2^{(n)}(Q^2)$ which are shown in Figure 2. At large Q^2 , far beyond the nodal region,

$$F_2^n(x, Q^2) \propto \left(\frac{x_0}{x}\right)^{\Delta_n} \left[\frac{1}{\alpha_S(Q^2)}\right]^{4/3\Delta_n}. \quad (12)$$

Since the relevant variable is a power of the inverse gauge coupling the nodes are spaced by 2-3 orders of magnitude in Q^2 -scale and only the first two of them are in the accessible range of Q^2 . The first nodes of $F_2^{(n)}(Q^2)$ are located at $Q^2 \sim 20 - 60 \text{ GeV}^2$. Hence, only the leading structure function contributes significantly in this region. This explains the precocious BFKL asymptotics found from the numerical solution of the BFKL equation [8].

In the accessible range of x the subleading contributions are numerically large. In particular, the BFKL expansion of the Born approximation dipole cross section which is used as the boundary condition at $x = x_0 = 0.03$ suggests more than 60% contribution from $F_2^{(n)}$ with $n > 0$. This implies the subleading terms determine the Q^2 dependence of $F_{2p}(Q^2)$ at $x = x_0$ and simultaneously the x dependence of the structure functions. Notice that in the pre-nodal region of $Q^2 \lesssim 20 \text{ GeV}^2$ the leading and subleading structure functions are very similar in shape. This explains the failure of the early GLDAP fits: only the limited region of $Q^2 \lesssim 10 \text{ GeV}^2$ was accessible and they




FIGURE 2. The modulus of structure function $F_2^{(n)}(Q^2)$ for $n = 0, 1, 2, 3$

could not catch and correctly describe a very different x -evolution of pomerons with different n .

At small x only the region $Q^2 \lesssim 10^3 \text{ GeV}^2$ is accessible. In this range the structure functions with $n \geq 3$ are hardly distinguishable. Besides, the splitting of the intercept with $n \geq 3$ is much smaller than for $n = 0, 1, 2$. Hence, the Regge expansion (2) can be truncated at $n = 2$ and $F_2^{(2)}(Q^2)$ comprises contributions from all poles with $n \geq 2$.

The BFKL equation allows one to determine the intercepts and structure functions $F_2^{(n)}(Q^2)$. The only adjustable parameters are "the pole residues" A_0, A_1, A_2 in (2) which are fixed, in fact, by the boundary condition at $x = x_0$. With the proper account of the valence [9] and non-perturbative [8] corrections to (2) we arrive at the three-pole-approximation which appears to be very successful when confronted with the data [10]. in a wide kinematical range (Figure 3). The effective pomeron intercept

$$\Delta_{eff} = -\frac{\partial \log F_{2p}(x, Q^2)}{\partial \log x} \quad (13)$$

gives an idea of the role of the subleading singularities. The intercept Δ_{eff} calculated with the experimental kinematic constraints is much smaller than $\Delta_0 = 0.4$ which is expected to dominate asymptotically. The agreement of our numerical estimates with the H1 determination (Figure 4) is quite satisfactory.

Acknowledgments I would like to thank José Repond for the kind hospitality in Chicago and careful reading the manuscript.




FIGURE 3. Three-Pole Approximation vs. H1, ZEUS and E665 data.

REFERENCES

1. Gribov V.N., and Lipatov L.N., *Sov. J. Nucl. Phys.* **15**, 438 (1972); Dokshitzer Yu.L., *Sov. Phys. JETP* **46**, 641 (1977); Altarelli G., and Parisi G., *Nucl. Phys.* **B126**, 297 (1977).
2. Fadin V.S., Kuraev E.A., and Lipatov L.N., *Sov. Phys. JETP* **44**, 443 (1976); *ibid* **45**, 199 (1977).
3. Lipatov L.N., *Sov. Phys. JETP* **63**, 904 (1986)
4. Nikolaev N.N., Zakharov B.G., and Zoller V.R., *JETP Letters* **59**, 8 (1994).
5. Nikolaev N.N., Zakharov B.G., *Phys. Letters* **B327**, 157 (1994).
6. Karl G., and Novikov V. *Phys. Rev* **D51**, 5069 (1995).
7. Nikolaev N.N., Zakharov B.G., and Zoller V.R., *JETP Letters* **60**, 678 (1994).
8. Nikolaev N.N., and Zakharov B.G., *Phys. Letters* **B333**, 250 (1994).
9. Glück M., Reya E., and Vogt A, *Z. Phys.* **C67**, 433 (1995).
10. H1 Collab., *Nucl. Phys.* **B439**, 471 (1995); ZEUS Collab., *Z. Phys.* **C69**, 607 (1996); E665 Collab., *Phys. Rev.* **D54**, 3006 (1996).




FIGURE 4. Effective intercept vs. H1 data.